



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2017

---

## **Phase I Dose-Escalation and -Expansion Study of the BRAF Inhibitor Encorafenib (LGX818) in Metastatic BRAF-Mutant Melanoma**

Delord, Jean-Pierre ; Robert, Caroline ; Nyakas, Marta ; McArthur, Grant A ; Kudchakar, Ragini ; Mahipal, Amit ; Yamada, Yasuhide ; Sullivan, Ryan ; Arance, Ana ; Kefford, Richard F ; Carlino, Matteo S ; Hidalgo, Manuel ; Gomez-Roca, Carlos ; Michel, Daniela ; Seroutou, Abdelkader ; Aslanis, Vassilios ; Caponigro, Giordano ; Stuart, Darrin D ; Moutouh-de Parseval, Laure ; Demuth, Tim ; Dummer, Reinhard

DOI: <https://doi.org/10.1158/1078-0432.CCR-16-2923>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-139238>

Journal Article

Accepted Version

Originally published at:

Delord, Jean-Pierre; Robert, Caroline; Nyakas, Marta; McArthur, Grant A; Kudchakar, Ragini; Mahipal, Amit; Yamada, Yasuhide; Sullivan, Ryan; Arance, Ana; Kefford, Richard F; Carlino, Matteo S; Hidalgo, Manuel; Gomez-Roca, Carlos; Michel, Daniela; Seroutou, Abdelkader; Aslanis, Vassilios; Caponigro, Giordano; Stuart, Darrin D; Moutouh-de Parseval, Laure; Demuth, Tim; Dummer, Reinhard (2017). Phase I Dose-Escalation and -Expansion Study of the BRAF Inhibitor Encorafenib (LGX818) in Metastatic BRAF-Mutant Melanoma. *Clinical Cancer Research*:1-38.

DOI: <https://doi.org/10.1158/1078-0432.CCR-16-2923>

## Phase I Dose-Escalation and -Expansion Study of the BRAF Inhibitor

### Encorafenib (LGX818) in Metastatic BRAF-Mutant Melanoma

Jean-Pierre Delord,<sup>1</sup> Caroline Robert,<sup>2</sup> Marta Nyakas,<sup>3</sup> Grant A. McArthur,<sup>4</sup> Ragini Kudchakar,<sup>5</sup> Amit Mahipal,<sup>6</sup> Yasuhide Yamada,<sup>7</sup> Ryan Sullivan,<sup>8</sup> Ana Arance,<sup>9</sup> Richard F. Kefford,<sup>10,11</sup> Matteo S. Carlino,<sup>10</sup> Manuel Hidalgo,<sup>12</sup> Carlos Gomez-Roca,<sup>1</sup> Daniela Michel,<sup>13</sup> Abdelkader Seroutou,<sup>13</sup> Vassilios Aslanis,<sup>13</sup> Giordano Caponigro,<sup>14</sup> Darrin D. Stuart,<sup>14</sup> Laure Moutouh-de Parseval,<sup>13</sup> Tim Demuth,<sup>13</sup> Reinhard Dummer<sup>15</sup>

<sup>1</sup>Institut Universitaire du Cancer, Toulouse, France; <sup>2</sup>Gustave Roussy and Paris-Sud University, Villejuif, France; <sup>3</sup>Oslo University Hospital, Norway; <sup>4</sup>Peter MacCallum Cancer Centre and the University of Melbourne, Australia; <sup>5</sup>Winship Cancer Institute of Emory University, Atlanta, GA, USA; <sup>6</sup>Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA; <sup>7</sup>National Cancer Center Hospital, Tokyo, Japan; <sup>8</sup>Massachusetts General Hospital, Boston, MA, USA; <sup>9</sup>Department of Medical Oncology and Translational Genomics and Targeted Therapeutics in Solid Tumors, Hospital Clínic, Barcelona, Spain; <sup>10</sup>Crown Princess Mary Cancer Centre Westmead Hospital, Melanoma Institute Australia, University of Sydney, Sydney, NSW, Australia; <sup>11</sup>Macquarie University, Sydney, NSW, Australia; <sup>12</sup>Spanish National Cancer Research Centre (CNIO) and Centro Integral Oncologico Clara Campal, Madrid, Spain; <sup>13</sup>Novartis Pharma AG, Basel, Switzerland; <sup>14</sup>Novartis Institutes for Biomedical Research, Cambridge, MA, USA; <sup>15</sup>University Hospital Zurich, Zurich, Switzerland

### Running head (65 characters max. incl. spaces)

Phase I study: encorafenib in *BRAF*-mutant melanoma

## **Key Words**

Melanoma, BRAF, encorafenib, binimetinib, Phase I

## **Corresponding Author: Professor Jean-Pierre Delord**

Clinical Research Unit, Institut Universitaire du Cancer, Oncopole, 1, avenue Irène  
Joliot-Curie, 31059, Toulouse-France

Tel: 330-561-42-41-14; Fax: 330-561-42-42-64; Email: [delord.jean-pierre@iuct-oncopole.fr](mailto:delord.jean-pierre@iuct-oncopole.fr)

## Translational Relevance

*BRAF* mutations occur in approximately 50% of melanoma cases. Treatment with currently approved BRAF inhibitors (BRAFi) in patients with *BRAF*-mutant melanoma results in disease progression within 6-8 months of starting treatment. Combined therapies of BRAF and MEK inhibitors (dabrafenib + trametinib; vemurafenib + cobimetinib) have been approved after demonstrating superior efficacy compared with BRAFi monotherapy. Encorafenib (LGX818), a selective BRAFi, was evaluated preclinically and in a single-agent phase I clinical study. Encorafenib showed more potent and prolonged pharmacodynamic activity compared with vemurafenib and dabrafenib. Clinical activity was observed in both BRAFi-pretreated and BRAFi-naïve patients at the recommended phase II dose (300 mg QD). The safety profile was distinct compared with other approved BRAFis, with lower rates of cutaneous squamous cell carcinoma and photosensitivity, but higher rates of Palmar-Plantar Erythrodysesthesia and facial paresis. These findings highlight the potential of encorafenib in the treatment of *BRAF*-mutant melanoma in future combination studies.

## Abstract

**Purpose:** Encorafenib, a selective BRAF inhibitor (BRAFi), has a pharmacological profile that is distinct from that of other clinically active BRAFis. We evaluated encorafenib in a phase I study in patients with BRAFi treatment-naïve and pretreated *BRAF*-mutant melanoma.

**Experimental Design:** The pharmacological activity of encorafenib was first characterized preclinically. Encorafenib monotherapy was then tested across a range of once-daily (QD; 50-700 mg) or twice-daily (75-150 mg) regimens in a phase I, open-label, dose-escalation and -expansion study in adult patients with histologically confirmed advanced/metastatic *BRAF*-mutant melanoma. Study objectives were to determine the maximum tolerated dose (MTD) and/or recommended phase II dose (RP2D), characterize the safety and tolerability and pharmacokinetic profile, and assess the preliminary antitumor activity of encorafenib.

**Results:** Preclinical data demonstrated that encorafenib inhibited BRAF V600E kinase activity with a prolonged off-rate and suppressed proliferation and tumor growth of *BRAF* V600E-mutant melanoma models. In the dose-escalation phase, 54 patients (29 BRAFi-pretreated and 25 BRAFi-naïve) were enrolled. Seven patients in the dose-determining set experienced dose-limiting toxicities. Encorafenib at a dose of 300 mg QD was declared the RP2D. In the expansion phase, the most common all-cause adverse events were nausea (66%), myalgia (63%), and palmar-plantar erythrodysesthesia (54%). In BRAFi-naïve patients, the overall response rate (ORR) and median progression-free survival (mPFS) were 60% and 12.4 months (95% CI, 7.4-NR). In BRAFi-pretreated patients, the ORR and mPFS were 22% and 1.9 months (95% CI, 0.9-3.7).

**Conclusions:** Once-daily dosing of single-agent encorafenib had a distinct tolerability profile and showed varying antitumor activity across BRAFi-pretreated and BRAFi-naïve patients with advanced/metastatic melanoma.

## INTRODUCTION

Melanoma is responsible for the majority of skin cancer deaths worldwide (1), and approximately 232,000 new cases of melanoma were diagnosed in 2012 (2). Most melanomas harbor alterations in components of the RAF/MEK/mitogen-activated protein kinase (MAPK) signaling transduction pathway, which drives cell proliferation, differentiation, and survival and can therefore lead to malignant cell proliferation (3).

Mutations in the *BRAF* gene occur in approximately 50% of melanoma cases (4). Over 90% of *BRAF* mutations are valine 600 substitutions (V600) (4). Vemurafenib and dabrafenib are currently the only approved BRAF inhibitors (BRAFi) for the treatment of *BRAF* V600-mutant melanoma (5-8). Unfortunately, most patients treated with BRAFi develop resistance and exhibit disease progression within 6-8 months of starting therapy, despite substantial initial tumor regression (9). Dual inhibition of the MAPK pathway with combined BRAF and MEK inhibitors have become the new standard of care in melanoma therapy following the approval of dabrafenib plus trametinib, and vemurafenib plus cobimetinib combination therapies, which demonstrated superior response to single-agent BRAFi in advanced melanoma (10-13).

BRAFi are associated with a number of proliferative and malignant cutaneous side effects, including cutaneous squamous cell carcinomas (SCC) and keratoacanthomas. Paradoxical activation of wild-type BRAF/CRAF is proposed to be the underlying mechanism of these adverse effects (AEs) (14). Highly selective BRAFi that are effective at low doses and with fewer side effects are ideal candidates for both single-agent and combination treatments. Encorafenib is a

potent, selective RAF kinase inhibitor with promising activity in preclinical models, including increased potency compared with vemurafenib (15). Presented here are results from preclinical studies demonstrating the pharmacologic properties of encorafenib as well as a phase I dose-escalation and dose-expansion study of single-agent encorafenib in patients with metastatic *BRAF*-mutant melanoma.

## METHODS

### *Pharmacologic characterization of encorafenib in preclinical models*

The potency of encorafenib, vemurafenib, and dabrafenib were measured in biochemical assays using methods previously described (16). Inhibition of BRAF V600E was evaluated in cell lines and xenograft tumors by measuring phosphorylated ERK (pERK) and total ERK using an immunoassay (Meso Scale Discovery). Cell proliferation assays were performed following 3 days of drug exposure with viability measured using CellTiter Glo 2.0. Mouse xenograft studies were performed in 6- to 8-week-old female nude mice following subcutaneous implantation of A375 cells ( $5 \times 10^6$ ) in 50% Matrigel or 3-mm  $\times$  3-mm pieces of the HMEX1906 primary human melanoma xenograft model. All laboratory animal work was conducted following appropriate laws and guidelines of the US Department of Agriculture, Office of Laboratory Animal Welfare, Association for Assessment and Accreditation of Laboratory Animal Care, and the Guide for the Care and Use of Laboratory Animals.



## *Phase I clinical trial*

### **Patient eligibility**

Key inclusion criteria were: age  $\geq 18$  years old; evaluable, histologically confirmed, locally advanced/metastatic *BRAF* V600-mutant melanoma (unresectable stage IIIB-IV); and World Health Organization performance status  $\leq 2$ . Key exclusion criteria were: prior MEK inhibitor therapy; prior systemic anticancer treatment  $\leq 4$  weeks, or major surgery  $\leq 2$  weeks prior to starting the study; symptomatic/untreated leptomeningeal disease or brain metastases; acute/chronic pancreatitis; clinically significant cardiac disease; or impairment of gastrointestinal function. All patients were required to provide written informed consent obtained before any screening procedure. This study was conducted according to the ethical principles of the Declaration of Helsinki.

### **Study design and treatments**

This phase I open-label, multicenter study included a dose-escalation and -expansion phase (NCT01436656). Patients were treated until disease progression, unacceptable toxicity, or withdrawal of informed consent. The primary objective was to determine the maximum tolerated dose (MTD) and/or recommended phase II dose (RP2D). Secondary objectives were to characterize the safety and tolerability and assess preliminary clinical antitumor activity and the pharmacokinetic profile of oral encorafenib. Exploratory objectives included assessing potential correlations of molecular status of RAF/MEK/MAPK and PI3K/AKT signaling molecules with clinical outcomes.

In the dose-escalation phase, patients were treated with either once-daily (QD) or twice-daily (BID) encorafenib in either microemulsion (50 or 100 mg QD) or capsule (50-700 mg QD, or 75-150 mg BID) formulation, administered in 28-day cycles. As the microemulsion and capsule formulations did not differ significantly in pharmacokinetic exposures or safety profiles at 50 and 100 mg QD, patients treated with the microemulsion were later transitioned to capsules at the same dose, for convenience. A Bayesian logistic regression model employing the escalation with overdose control (EWOC) principle was used to guide dose escalation and to estimate the MTD. The study started treating cohorts of 1–3 patients at the prespecified dose levels. Cohorts were expanded between 3–6 patients when dose-limiting toxicities (DLTs) were identified, or additional cohorts of 1–6 evaluable patients were added when a dose level showed signs of activity and satisfied EWOC criteria. The dose-escalation phase was continued until the MTD and/or RP2D of oral encorafenib was determined. The maximum tolerated dose was defined as the highest drug dose expected, based on the Bayesian logistic regression model, to not cause a DLT in more than 33% of the treated patients in the first cycle of encorafenib treatment

The dose-expansion phase evaluated the safety and preliminary antitumor activity at the estimated MTD/RP2D in patients with locally advanced/metastatic melanoma (BRAFi-naive or BRAFi-pretreated). Most patients enrolled received either 300 or 450 mg QD encorafenib. In addition, a small cohort of patients was treated with stepwise dosing of encorafenib from 300 mg (up to day [D] 15) to 450 mg (from D15 onward if patients experienced no clinically significant grade 3/4 AEs at the lower

dose level). A third group of patients with metastatic colorectal cancer were enrolled in this trial (17) but their data will be published in a separate manuscript.

## **Study procedures**

**Safety assessments:** Data regarding all adverse events (AEs) and serious AEs (SAEs) were collected, along with the severity and relationship to the study drug; AEs and SAEs were graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) v4.0 (18).

**Efficacy assessments:** All potential sites of tumor lesions were locally assessed by radiological techniques every 8 weeks per Response Evaluation Criteria In Solid Tumors (RECIST) v1.0 (19) for the dose-escalation phase and per RECIST v1.1 (20) for the dose-expansion phase. All complete and partial responses were confirmed by a second assessment at least 4 weeks later.

**Pharmacokinetic assessments:** Blood samples for full pharmacokinetic profiles were collected on cycle 1 day 1 (C1D1), C1D8, and C1D15. Patients treated with stepwise dosing had samples collected on C1D1, C1D15, and C2D1 for dose expansion.

Plasma encorafenib concentrations were measured using a validated liquid chromatography–tandem mass spectrometry assay.

**Somatic mutation and biomarker assessments:** Representative tumor tissue was used to determine the mutation status of other genes relevant to RAF/MEK/ERK signaling to identify potential predictive markers of efficacy. Next-generation sequencing was conducted at Foundation Medicine using a T5a panel.

## Statistical analysis

The analysis was based on all patients' data from the dose-escalation and dose-expansion phases up to the time when all patients had completed at least 8 cycles of treatment or discontinued the study. Data are summarized with respect to demographic and baseline characteristics and efficacy and safety observations. Continuous variables are summarized as median and ranges. Categorical variables are expressed as frequencies and percentages. Progression-free survival (PFS) and overall survival (OS) were estimated using the Kaplan-Meier method and medians were reported with corresponding 95% CIs.

## RESULTS

### Preclinical Characterization

#### *Pharmacologic characterization of encorafenib in preclinical models*

Encorafenib inhibited BRAF V600E kinase activity in a biochemical assay at similar concentrations as dabrafenib and vemurafenib (Supplementary Table S1); however, the dissociation half-life was considerably increased (>30 hours) compared with dabrafenib (2 hours) and vemurafenib (0.5 hours; Supplementary Table S1 and Figure 1A). Washout experiments in A375 cells confirmed the long dissociation half-life of encorafenib, as the phosphorylated ERK half maximal effective concentration (pERK EC<sub>50</sub>) only shifted 2-fold following washout compared with 14-fold and 23-fold for dabrafenib and vemurafenib, respectively (Supplementary Table S1). In a broader survey of *BRAF* V600-mutant cell lines, encorafenib was more potent at inhibiting proliferation, with most cell lines having IC<sub>50</sub> <40 nM, while slightly higher concentrations of dabrafenib (<100 nM) and significantly higher concentrations of

vemurafenib (<1  $\mu$ M) were required to inhibit proliferation of most cell lines (Supplemental Figure S1).

Single-dose pharmacokinetic/pharmacodynamic studies in human melanoma xenograft models (*BRAF* V600E) showed that oral doses of encorafenib as low as 6 mg/kg resulted in strong (75%) and sustained (>24 h) decreases in pERK (Figure 1B), consistent with a prolonged dissociation half-life. Encorafenib induced tumor regression in multiple *BRAF*-mutant human tumor xenograft models grown in immune-compromised mice (Figures 1C and 1D). Although maximum encorafenib antitumor activity appeared to be achieved at 5 mg/kg BID (Figure 1E), higher doses were required to prevent the emergence of resistance and improve conditional survival (Figure 1F).

Encorafenib was well tolerated, and exposure was dose proportional at all doses tested in mice (Supplemental Figure S2A, B). In order to determine the effect of encorafenib on cell proliferation that may potentially lead to SSC growth, epithelial hyperplasia and hyperkeratosis in the forestomach of mice were investigated over a dose range. Epithelial hyperplasia and hyperkeratosis were observed at very high encorafenib doses, in contrast to both vemurafenib (4/5 mice) and dabrafenib (5/5 mice), with which gastric hyperplasia and hyperkeratosis were observed at clinically relevant doses (Supplementary Figure S2C).

## Phase I Clinical Trial

### *Patient disposition and characteristics*

Fifty-four patients with *BRAF*-mutant melanoma (25 BRAFi-naïve and 29 BRAFi-pretreated patients) were enrolled and treated in the dose-escalation phase. At the time of data analysis cutoff (18 August 2014), 49 patients had discontinued treatment (22 BRAFi-naïve and 27 BRAFi-pretreated patients); the most common reason for treatment discontinuation was disease progression (n=41; 16 [64.0%] BRAFi-naïve and 25 [86.2%] BRAFi-pretreated patients, respectively), followed by AEs (n=5; 9.3%). Thirty-five additional patients (17 BRAFi-naïve and 18 pretreated patients) were treated in the dose-expansion phase; 27 of these patients discontinued treatment by the time of data cutoff (11 BRAFi-naïve and 16 BRAFi-pretreated patients). The most common reason for discontinuation was disease progression (n=17; 3 [17.6%] BRAFi-naïve and 14 [77.8%] BRAFi-pretreated patients, respectively), followed by AEs (n=8; 22.9%). In BRAFi-pretreated patients, the median time from prior BRAFi treatment to the start of encorafenib was 38.5 days (range, 11-331). Patient characteristics are summarized in Table 1.

### *Determination of MTD/RP2D*

The dose-determining set consisted of 49 patients treated across 11 dose levels. Seven patients (14.3%), of whom 3 were treated above 450 mg QD, experienced at least 1 DLT during cycle 1. The most frequent DLT was neuralgia (2 patients, 4.1%). All other DLTs occurred in 1 patient each (asthenia, confusional state, diarrhea, facial paresis, fatigue, headache, insomnia, musculoskeletal pain, neck pain, palmar-plantar erythrodysesthesia [PPED] syndrome and rash; 2% each; Supplementary Table S2). Based on the Bayesian model, 450 mg QD was declared the MTD.

In the dose-expansion phase, 9 patients (33.3%), of whom 6 were treated at the 450 mg QD MTD, experienced at least 1 DLT during cycle 1. At the MTD, DLTs included myalgia (33.3%), arthralgia (26.7%), fatigue (20.0%), asthenia (13.3%), seventh nerve paralysis (6.7%), and insomnia (6.7%; Supplementary Table S2). All DLTs were grade 3. Of these 9 patients, 7 required a dose reduction to 300 mg QD. At this time, the 450 mg QD MTD no longer met the EWOC criteria with a risk of excessive toxicity of 37.9%. Thus, the 300 mg QD dose level was better tolerated and declared the RP2D. The risk of excessive toxicity at 300 mg QD was 0.6%.

Review of the BID dosing cohort revealed that the tolerability of this dosing schedule did not justify BID dosing in view of potentially reduced compliance compared with the QD dosing schedule. In conjunction with the known long dissociation half-life of encorafenib, a QD dosing regimen was selected for future dosing.

### *Safety and tolerability*

#### **Dose-escalation phase**

The median duration of exposure was 15.1 weeks (range, 0.7-142), and 5 patients had ongoing treatment at the time of data cutoff. All 54 patients experienced at least 1 AE (Table 2). Grade 3/4 AEs were reported in 38 patients (70.4%); 6 patients (11.1%) had grade 4 AEs. The most common AEs (>40%) were PPED syndrome (51.9%), hyperkeratosis (50.0%), arthralgia (44.4%), nausea (44.4%), and pruritus (40.7%). Fifty-two patients had at least 1 AE suspected to be study drug related; of these, 18 patients (33.3%) had grade 3 AEs. Grade 3/4 study drug-related AEs reported (≥5% of patients) were fatigue (9.3%) and PPED syndrome (7.4%). Notably, the incidence of secondary neoplasms was low; SCC and keratoacanthoma were reported in 2 patients (3.7%) each. Three patients (5.6%) experienced facial paresis

(including facial paresis and seventh nerve paralysis) at 100-mg QD, 100-mg BID, and 450-mg QD dose levels. Pyrexia was reported in 2 patients (3.7%) and photosensitivity reactions were reported in 7 patients (13%).

Five patients (9.3%) experienced AEs leading to treatment discontinuation; these were PPED syndrome (3.7%, 100 and 450 mg QD), headache (1.9%, 700 mg QD), hyperkeratosis (1.9%, 550 mg QD), and neuralgia (1.9%, 300 mg QD). Eleven on-treatment deaths (defined as occurring during treatment or within 30 days of the last dose of study medication) were reported. One death, which was due to general health deterioration, was considered to be related to the study treatment by the study investigator; however, the role of disease progression, which was documented 19 days prior to death, may also be a plausible explanation for the outcome. The patient had ongoing events at the time of death, including dysphagia, confusional state, and a new melanoma. The remaining 10 on-treatment deaths (1 during treatment and 9 within 30 days of the last encorafenib dose) were due to progressive disease and unrelated to the study treatment.

### **Dose-expansion phase**

The median duration of exposure for all patients was 12.1 weeks (range, 0.1-90.7), and 8 patients had treatment ongoing at the time of data cutoff. In patients who had been pretreated with a BRAFi, exposure duration was 8.9 weeks (range, 0.4-90.7) in the MTD group and 5.6 weeks (range, 0.4-10.9) in the RP2D group. For BRAFi-naïve patients this was 6.3 weeks (range, 0.1-71.7) in the MTD group and 38.6 weeks (range, 0.4-53.9) in the RP2D group.



AEs experienced in the dose-expansion phase were similar to those experienced in the dose-escalation phase (Table 2). Thirty-four (97.1%) patients had at least 1 AE suspected to be study drug-related. The most frequent study drug-related AEs (>40%) were myalgia (62.9%), nausea (57.1%), PPED syndrome (54.3%), arthralgia (48.6%), vomiting (45.7%), alopecia (42.9%), dry skin (42.9%), and insomnia (40.0%). One patient (2.9%) experienced SCC of the head and neck, and keratoacanthoma was not observed. Four patients (11.4%) experienced facial paresis (seventh nerve paralysis) at 300 mg QD (n=1), at 450 mg QD (n=2), and in the stepwise group (n=1). Pyrexia was reported in 2 patients (5.7%), and photosensitivity reactions were reported in 1 patient (2.9%).

Eight patients (22.9%) had AEs leading to discontinuation of study drug; the most common AEs were arthralgia and myalgia (5.7% each). Two on-treatment deaths (both within 30 days of the last dose of encorafenib) were reported and were not considered to be related to study treatment.

### *Pharmacokinetics*

Encorafenib was rapidly absorbed and detectable in plasma at 0.5 h postdose and across all dose levels peaked ( $T_{max}$ ) at approximately 2 h (Figure 2). Terminal half-life ( $T_{1/2}$ ) was short (2.9-4.4 h), remained constant across doses, and was similar between C1D1 and C1D15 (Supplementary Table S3). Plasma concentrations of encorafenib were lower on C1D15 than C1D1 across all dose levels, suggesting time-dependent pharmacokinetics. Plasma clearance was increased 2-fold between C1D1 and C1D15, while the area under the plasma concentration-time curve for a dosing interval ( $AUC_T$ ) and maximum plasma concentration ( $C_{max}$ ) showed that systemic exposure was also lower at C1D15 (Supplementary Table S3), possibly

due to autoinduction via the CYP3A4 pathway. Trough plasma concentrations did not show an appreciable difference by C2D1, suggesting that steady-state conditions had been reached by C1D15. Mixed-effect modeling on QD capsule data of C1D1 and C1D15 confirmed that AUC and  $C_{\max}$  were approximately dose proportional over the complete dose range (Supplementary Table S3).

### *Efficacy*

In the dose-escalation phase, the overall response rates (ORRs) for BRAFi-naïve and BRAFi-pretreated patients were 60.0% and 10.3%, respectively (Table 3, Figure 3A). In the dose-expansion phase, the ORRs were 60.0% and 22.2% for BRAFi-naïve and BRAFi-pretreated patients, respectively (Table 3; Figure 3B). PFS and OS were assessed for patients in the dose-expansion phase only. Median PFS for BRAFi-naïve patients was 12.4 months (95% CI, 7.4-NR), and the PFS rate at 6 months was 91.7% (95% CI, 76.0-100). Median PFS for BRAFi-pretreated patients was 1.9 months (95% CI, 0.9-3.7), with a PFS rate at 6 months of 11.8% (95% CI, 0.0-27.1; Figure 3C). Median OS was not reached for BRAFi-naïve patients and was 9.07 months (95% CI, 3.68-10.84) for BRAFi-pretreated patients (Figure 3D).

### *Biomarker analysis*

The mutation status of archival or fresh tumor samples collected at baseline was analyzed centrally by next-generation sequencing. No correlation was observed between the best percentage change from baseline and mutation status (Supplementary Figure S3A and B).

## DISCUSSION

Preclinical studies demonstrate that encorafenib is more potent and has prolonged pharmacodynamic activity compared with approved BRAF inhibitors vemurafenib and dabrafenib. Antitumor activity of encorafenib was demonstrated in human *BRAF* V600E-mutant melanoma xenografts at doses as low as 5 mg/kg; however, higher doses were required to prevent the emergence of resistance in this model. Positive preclinical findings led to further investigation of encorafenib in clinical trials. Data from a more comprehensive investigation of the preclinical comparative pharmacology of encorafenib and other BRAF inhibitors are in preparation for publication.

The primary objective of this phase I study was to determine the MTD and/or RP2D of encorafenib. The MTD was established at 450 mg QD based on data from the dose-escalation phase; however, after reviewing the safety and tolerability data in the dose-expansion phase, the MTD was not considered to be a viable RP2D because 7 of 9 patients who exhibited a DLT during the first cycle required a dose reduction to 300 mg QD. Therefore, the 300-mg QD dose level was declared the RP2D. Review of the overall safety profile of the 450-mg QD and 300-mg QD doses continued to show meaningfully higher rates of DLTs and grade 3 AEs at the MTD, supporting 300 mg QD as the RP2D for encorafenib as a single agent.

Encorafenib showed signs of promising activity in this study, with ORRs in the dose-expansion phase of 60% in patients with melanoma who were BRAFi-naïve. Minimal efficacy was seen in patients who received prior BRAFi treatment. BRAFi-pretreated patients received encorafenib between 1.5-47 weeks following their BRAFi

treatment. Objective responses to BRAFi retreatment have been reported in patients progressing on BRAFi therapy following a treatment break (21, 22). Therefore, treatment interruption may account for some of the responses observed in pretreated patients in this study due to removal of the selective pressure of MAPK inhibition. Correlative analyses between the length of the BRAFi treatment-break interval and response to subsequent BRAFi will help further understand tumor response dynamics. Encorafenib showed durable response in BRAFi-naïve patients, with a median PFS of 12.4 months—longer than that reported in vemurafenib (5.3 months) (23) and dabrafenib (4.5-6.2 months) in the same setting (24). Data from the randomized COLUMBUS trial confirm this observation (25).

Encorafenib was well tolerated, with most AEs being grade 2 or lower in severity. Single-agent BRAFis have been associated with characteristic AEs, which are proposed to be the result of aberrant wild-type BRAF/CRAF activation. Dermatologic events were among the most common AEs observed in the dose-escalation and -expansion phases. PPED and hyperkeratosis were reported in over 40% of patients, higher than the rates reported with the approved BRAF inhibitors. Although hyperkeratosis is a characteristic and diagnostic feature of PPED according to the NCI CTCAE v4.0 reporting criteria (18), hyperkeratosis was often experienced in locations other than the hands and feet. Cutaneous SCC has been reported in 12%-21% of patients with melanoma following treatment with vemurafenib monotherapy (23, 26, 27) and in 9%-19% of patients following dabrafenib monotherapy (11, 24, 28), in contrast to the low rates (3%-4%) observed in this study. These observations are intriguing given the lower rate of hyperplasia observed in the gastric epithelium of mice treated with encorafenib. A recent study

profiling the paradoxical activation of ERK across the BRAF inhibitors vemurafenib, dabrafenib, and encorafenib suggests that differences in the potency of pERK induction in *HRAS<sup>mut</sup>*, *BRAF<sup>wt</sup>* keratinocytes relative to the potency of pERK inhibition in *BRAF* V600E melanoma cells may account for the varying rates of cutaneous SCC between the compounds (29).

Facial paresis has been reported in cases of patients treated with vemurafenib (30, 31). Symptoms typically resolve when treatment is stopped and may be resolved with steroid treatment, suggesting a possible autoimmune mechanistic trigger. Facial paresis (regardless of causality) was observed in 7 patients (7.9%) treated with encorafenib. The low rates of pyrexia, photosensitivity, and cutaneous SCC suggest a tolerability profile that is different from either vemurafenib or dabrafenib (5, 7). Further studies are needed to better understand the mechanisms that underlie these differences, which may include on-target and off-target effects (32).

It is important to note that the standard of care with BRAF-targeted therapy has changed dramatically since the commencement of this study. Specifically, there are now 3 randomized trials demonstrating superior response rates, PFS, and OS in patients receiving combinations of BRAF and MEK inhibitors (dabrafenib plus trametinib, and vemurafenib plus cobimetinib) compared with single-agent BRAFi therapy (11-13, 28), which led to the approval of these combinations for the treatment of patients with unresectable or metastatic melanoma with a *BRAF* V600E or V600K mutation. Preliminary results from combination studies have suggested that pairing encorafenib with the MEK inhibitor binimetinib mitigates some dermatologic AEs that were associated with single-agent encorafenib, a finding that

has also been reported with the combination of dabrafenib and trametinib (33), and appears to be associated with efficacy comparable to other BRAF/MEK inhibitor combinations (28, 34, 35). The efficacy and safety of encorafenib is currently being further investigated in combination with binimetinib, and compared with that of encorafenib or vemurafenib alone, in an ongoing phase III study (COLUMBUS, NCT01909453) (25).

### **Acknowledgments:**

The authors thank Fernando Salangsang, Kimberly Aardalen, Edward Lorenzana, Hanne Merritt, Susan Kaufman, and Majid Ghoddusi for generation of preclinical experiments. Financial support for medical editorial assistance was provided by Novartis Pharmaceuticals. We thank Zoe Crossman and Maria Alfaradhi for medical editorial assistance with this manuscript.

## References

1. American Cancer Society: Melanoma skin cancer.  
<http://www.cancer.org/cancer/skincancer-melanoma/index>
2. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, et al. Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-86.
3. Chappell WH, Steelman LS, Long JM, Kempf RC, Abrams SL, Franklin RA, et al. Ras/Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR inhibitors: Rationale and importance to inhibiting these pathways in human health. *Oncotarget* 2011;2:135-64.
4. Ascierto PA, Kirkwood JM, Grob JJ, Simeone E, Grimaldi AM, Maio M, et al. The role of BRAF V600 mutation in melanoma. *J Transl Med* 2012;10:85,5876-10-85.
5. Zelboraf (vemurafenib) [package insert]. South San Francisco, CA, Genentech USA, Inc; 2014.  
[http://www.accessdata.fda.gov/drugsatfda\\_docs/label/2014/202429s006lbl.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/202429s006lbl.pdf).
6. Zelboraf (vemurafenib) [summary of product characteristics]. Welwyn Garden City, UK: Roche; 2012. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Product\\_Information/human/002409/WC500124317.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002409/WC500124317.pdf).
7. Tafenlar (dabrafenib) [package insert]. Research Triangle Park, NC: GlaxoSmithKline; 2014.  
[http://www.accessdata.fda.gov/drugsatfda\\_docs/label/2014/202806s002lbl.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/202806s002lbl.pdf).

8. Tafenlar (dabrafenib) [summary of product characteristics]. Camberly, UK: Novartis; 2014. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/EPAR\\_-\\_Product\\_Information/human/002604/WC500149671.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002604/WC500149671.pdf).
9. Sullivan RJ, Flaherty KT. Resistance to BRAF-targeted therapy in melanoma. *Eur J Cancer* 2013;49:1297-304.
10. Robert C, Karaszewska B, Schachter J, Rutkowski P, Mackiewicz A, Stroiakovski D, et al. Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med* 2015;372:30-9.
11. Long GV, Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, et al. Dabrafenib and trametinib versus dabrafenib and placebo for Val600 BRAF-mutant melanoma: A multicentre, double-blind, phase 3 randomised controlled trial. *Lancet* 2015;386:444-51.
12. Larkin J, Ascierto PA, Dreno B, Atkinson V, Liskay G, Maio M, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med* 2014;371:1867-76.
13. Larkin J, Yan Y, McArthur GA, Ascierto PA, Liskay G, Maio M, et al. Update of progression-free survival (PFS) and correlative biomarker analysis from coBRIM: Phase III study of cobimetinib (cobi) plus vemurafenib (vem) in advanced *BRAF*-mutated melanoma. *J Clin Oncol* 2015;33(suppl; abstr 9006).
14. Holderfield M, Nagel TE, Stuart DD. Mechanism and consequences of RAF kinase activation by small-molecule inhibitors. *Br J Cancer* 2014;111:640-5.



15. Stuart DD, Li N, Poon DJ, Aardalen K, Kaufman S, Merritt H, et al. Preclinical profile of LGX818: A potent and selective RAF kinase inhibitor. *Cancer Res.* 2012;72:Abstract 3790.
16. Holderfield M, Merritt H, Chan J, Wallroth M, Tandeske L, Zhai H, et al. RAF inhibitors activate the MAPK pathway by relieving inhibitory autophosphorylation. *Cancer Cell* 2013;23:594-602.
17. Gomez-Roca CA, Delord J, Robert C, Hidalgo M, von Moos R, Arance A, et al. Encorafenib (LGX818), an oral BRAF inhibitor, in patients (pts) with BRAF V600E metastatic colorectal cancer (MCRC): Results of dose expansion in an open-label, phase 1 study. *Annals of Oncology* 2014;25(suppl 4):535P.
18. NCI. common terminology criteria for adverse events (CTCAE). Version 4.0. [http://evs.nci.nih.gov/ftp1/CTCAE/CTCAE\\_4.03\\_2010-06-14\\_QuickReference\\_5x7.pdf](http://evs.nci.nih.gov/ftp1/CTCAE/CTCAE_4.03_2010-06-14_QuickReference_5x7.pdf).
19. Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000;92:205-16.
20. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: Revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228-47.

21. Seghers AC, Wilgenhof S, Lebbe C, Neyns B. Successful rechallenge in two patients with BRAF-V600-mutant melanoma who experienced previous progression during treatment with a selective BRAF inhibitor. *Melanoma Res* 2012;22:466-72.
22. Mackiewicz-Wysocka M, Krokowicz L, Kocur J, Mackiewicz J. Resistance to vemurafenib can be reversible after treatment interruption: A case report of a metastatic melanoma patient. *Medicine (Baltimore)* 2014;93:e157.
23. Chapman PB, Hauschild A, Robert C, Haanen JB, Ascierto P, Larkin J, et al. Improved survival with vemurafenib in melanoma with BRAF V600E mutation. *N Engl J Med* 2011;364:2507-16.
24. Ascierto PA, Minor D, Ribas A, Lebbe C, O'Hagan A, Arya N, et al. Phase II trial (BREAK-2) of the BRAF inhibitor dabrafenib (GSK2118436) in patients with metastatic melanoma. *J Clin Oncol* 2013;31:3205-11.
25. Dummer R, Ascierto PA, Gogas HJ, Arance A, Mandala M, Liszkay G, et al. Results of COLUMBUS part 1: A phase 3 trial of encorafenib (ENCO) plus binimetinib (BINI) versus vemurafenib (VEM) or ENCO in BRAF-mutant melanoma. Society for Melanoma Research 2016;Boston, MA, US.
26. McArthur GA, Chapman PB, Robert C, Larkin J, Haanen JB, Dummer R, et al. Safety and efficacy of vemurafenib in BRAF(V600E) and BRAF(V600K) mutation-positive melanoma (BRIM-3): Extended follow-up of a phase 3, randomised, open-label study. *Lancet Oncol* 2014;15:323-32.

27. Flaherty KT, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, et al.  
 Inhibition of mutated, activated BRAF in metastatic melanoma. *N Engl J Med*  
 2010;363:809-19.
  
28. Flaherty KT, Infante JR, Daud A, Gonzalez R, Kefford RF, Sosman J, et al.  
 Combined BRAF and MEK inhibition in melanoma with BRAF V600 mutations. *N*  
*Engl J Med* 2012;367:1694-703.
  
29. Adelman CH, Ching G, Du L, Saporito RC, Bansal V, Pence LJ, et al.  
 Comparative profiles of BRAF inhibitors: The paradox index as a predictor of clinical  
 toxicity. *Oncotarget* 2016;7:30453-60.
  
30. Klein O, Ribas A, Chmielowski B, Walker G, Clements A, Long GV, et al. Facial  
 palsy as a side effect of vemurafenib treatment in patients with metastatic  
 melanoma. *J Clin Oncol* 2013;31:e215-7.
  
31. Shailesh FN, Singh M, Tiwari U, Hutchins LF. Vemurafenib-induced bilateral  
 facial palsy. *J Postgrad Med* 2014;60:187-8.
  
32. Vin H, Ojeda SS, Ching G, Leung ML, Chitsazzadeh V, Dwyer DW, et al. BRAF  
 inhibitors suppress apoptosis through off-target inhibition of JNK signaling. *Elife*  
 2013;2:e00969.
  
33. Long GV, Stroyakovskiy D, Gogas H, Levchenko E, de Braud F, Larkin J, et al.  
 Combined BRAF and MEK inhibition versus BRAF inhibition alone in melanoma. *N*  
*Engl J Med* 2014;371:1877-88.
  
34. Kefford R, Miller WH, Tan DS, Sullivan RJ, Long G, Dienstmann R, et al.  
 Preliminary results from a phase Ib/II, open-label, dose-escalation study of the oral

BRAF inhibitor LGX818 in combination with the oral MEK 1/2 inhibitor MEK162 in *BRAF* V600-dependent advanced solid tumors. J Clin Oncol 2013;31:(suppl; abstr 9029).

35. Sullivan RJ, Weber JS, Patel SP, Dummer R, Miller WH, Cosgrove D, et al. A phase Ib/II study of BRAF inhibitor (BRAFi) encorafenib (ENCO) plus MEK inhibitor (MEKi) binimetinib (BINI) in cutaneous melanoma patients naive to BRAFi treatment. J Clin Oncol 2015; 33:(suppl; abstr 9007).

**Table 1.** Baseline patient and disease characteristics

Characteristic	Dose Escalation		Dose Expansion		
	BRAF <sup>i</sup> Naive (n=25)	BRAF <sup>i</sup> Pretreated (n=29)	BRAF <sup>i</sup> Naive (n=15)	BRAF <sup>i</sup> Pretreated (n=18)	All Melanoma* (N=35)
Median age (range), years	51.0 (33-70)	50.0 (22-78)	61.0 (43-79)	54.0 (26-74)	59.0 (26-79)
Sex, n (%)					
Female	8 (32)	10 (35)	5 (33)	14 (78)	20 (57)
Male	17 (68)	19 (65)	10 (67)	4 (22)	15 (43)
WHO performance status, n (%)					
0	19 (76)	16 (55)	12 (80)	8 (44)	20 (57)
1	5 (20)	12 (41)	3 (20)	10 (56)	15 (43)
2	1 (4)	1 (3)	0 (0)	0 (0)	0 (0)
<i>BRAF</i> V600 mutation status, n (%)					
V600E	23 (92)	26 (90)	13 (87)	17 (94)	31 (89)
Other	2 (8)	3 (10)	2 (13)	1 (6)	4 (11)
LDH at baseline, n (%)					
≤ULN	16 (64)	16 (55)	10 (67)	4 (22)	14 (40)
>ULN	9 (36)	13 (45)	5 (33)	14 (78)	21 (60)
Stage at current diagnosis, n (%)					
Stage IIIb	1 (4)	0 (0)	1 (7)	0 (0)	1 (3)
Stage IIIc	0 (0)	0 (0)	1 (7)	0 (0)	1 (3)
Stage Iva	2 (8)	1 (3)	2 (13)	0 (0)	2 (6)
Stage IVb	0 (0)	4 (14)	4 (27)	1 (6)	5 (14)
Stage IVc	22 (88)	24 (83)	7 (47)	17 (94)	26 (74)
Previous BRAF <sup>i</sup> treatment, n (%)		29 (100)		18 (100)	
Vemurafenib		28 (97)		16 (89)	
Dabrafenib		0 (0)		2 (11)	
Investigational		1 (3)		0 (0)	
BRAF <sup>i</sup> Median time between discontinuation of BRAF <sup>i</sup> and study start, days (range)		26 (12-319)		38.5 (11-331)	



**Table 2.** Adverse events occurring in ≥30% of patients in the dose-expansion phase

Adverse Event	BRAFi Naive (n=15), n (%)		BRAFi Pretreated (n=18), n (%)		All Melanoma* (N=35), n (%)	
	All Grades	Grade 3/4	All Grades	Grade 3/4	All Grades	Grade 3/4
<b>Irrespective of study drug relationship</b>						
Nausea	9 (60.0)	1 (6.7)	13 (72.2)	3 (16.7)	23 (65.7)	4 (11.4)
Myalgia	8 (53.3)	2 (13.3)	13 (72.2)	3 (16.7)	22 (62.9)	5 (14.3)
PPED syndrome	6 (40.0)	0 (0.0)	11 (61.1)	3 (16.7)	19 (54.3)	4 (11.4)
Arthralgia	6 (40.0)	3 (20.0)	9 (50.0)	3 (16.7)	17 (48.6)	7 (20.0)
Vomiting	5 (33.3)	1 (6.7)	10 (55.6)	2 (11.1)	17 (48.6)	3 (8.6)
Insomnia	4 (26.7)	1 (6.7)	10 (55.6)	1 (5.6)	16 (45.7)	2 (5.7)
Alopecia	7 (46.7)	0 (0)	8 (44.4)	0 (0)	15 (42.9)	0 (0)
Dry skin	6 (40.0)	0 (0.0)	9 (50.0)	1 (5.6)	15 (42.9)	1 (2.9)
Hyperkeratosis	6 (40.0)	1 (6.7)	6 (33.3)	0 (0)	14 (40.0)	1 (2.9)
Headache	4 (26.7)	0 (0)	9 (50.0)	2 (11.1)	13 (37.1)	2 (5.7)
Keratosis pilaris	3 (20.0)	0 (0)	9 (50.0)	2 (11.1)	13 (37.1)	2 (5.7)
Pruritus	5 (33.3)	0 (0)	6 (33.3)	1 (5.6)	13 (37.1)	1 (2.9)
Asthenia	1 (6.7)	0 (0)	9 (50.0)	3 (16.7)	11 (31.4)	3 (8.6)
Constipation	5 (33.3)	0 (0)	5 (27.8)	0 (0)	11 (31.4)	0 (0)
<b>Suspected study drug-related</b>						
Myalgia	8 (53.3)	2 (13.3)	13 (72.2)	3 (16.7)	22 (62.9)	5 (14.3)
Nausea	8 (53.3)	1 (6.7)	11 (61.1)	1 (5.6)	20 (57.1)	2 (5.7)
PPED syndrome	6 (40.0)	0 (0)	11 (61.1)	3 (16.7)	19 (54.3)	4 (11.4)
Arthralgia	6 (40.0)	3 (20.0)	9 (50.0)	3 (16.7)	17 (48.6)	7 (20.0)
Vomiting	5 (33.3)	1 (6.7)	9 (50.0)	2 (11.1)	16 (45.7)	3 (8.6)
Alopecia	7 (46.7)	0 (0)	8 (44.4)	0 (0)	15 (42.9)	0 (0)
Dry skin	6 (40.0)	0 (0)	9 (50.0)	1 (5.6)	15 (42.9)	1 (2.9)
Insomnia	4 (26.7)	1 (6.7)	10 (55.6)	1 (5.6)	14 (40.0)	2 (5.7)
Hyperkeratosis	6 (40.0)	1 (6.7)	5 (27.8)	0 (0)	13 (37.1)	1 (2.9)
Keratosis pilaris	3 (20.0)	0 (0)	9 (50.0)	2 (11.1)	13 (37.1)	2 (5.7)
Pruritus	5 (33.3)	0 (0)	5 (27.8)	1 (5.6)	12 (34.3)	1 (2.9)
Asthenia	1 (6.7)	0 (0)	9 (50.0)	3 (16.7)	11 (31.4)	3 (8.6)

**Table 3.** Summary of best overall response for patients in the dose-escalation and dose-expansion phases

<b>Best Overall Response</b>	<b>Dose Escalation</b>		<b>Dose Expansion</b>		
	BRAFi	BRAFi	BRAFi	BRAFi	All
	Naive	Pretreated	Naive	Pretreated	Melanoma*
	(n=25), n (%)	(n=29), n (%)	(n=15), n (%)	(n=18), n (%)	(N=35), n (%)
CR	2 (8.0)	1 (3.4)	1 (6.7)	0	1 (2.9)
PR	13 (52.0)	2 (6.9)	8 (53.3)	4 (22.2)	13 (37.1)
Unconfirmed CR <sup>†</sup>	0	0	1 (6.7)	0	1 (2.9)
Stable disease	5 (20.0)	10 (34.5)	1 (6.7)	2 (11.1)	4 (11.4)
Unconfirmed	1 (4.0)	0	0	1 (5.6)	1 (2.9)
CR/PR <sup>‡</sup>					
Progressive disease	0	14 (48.3)	1 (6.7)	10 (55.6)	11 (31.4)
Unconfirmed	0	0	0	0	0
CR/PR					
Unknown	5 (20.0)	2 (6.9)	4 (26.7)	2 (11.1)	6 (17.1)
Unconfirmed	1 (4.0)	0	1 (6.7)	0	1 (2.9)
CR/PR <sup>§</sup>					
<b>ORR</b>	<b>15 (60.0)</b>	<b>3 (10.3)</b>	<b>9 (60.0)</b>	<b>4 (22.2)</b>	<b>14 (40.0)</b>
<b>95% CI</b>	<b>38.7-78.9</b>	<b>2.2-27.4</b>	<b>32.3-83.7</b>	<b>6.4-47.6</b>	<b>23.9-57.9</b>
Disease control rate	20 (80.0)	13 (44.8)	10 (66.7)	6 (33.3)	18 (51.4)
95% CI	59.3-93.2	26.4-64.3	38.4-88.2	13.3-59.0	34.0-68.6



## Table Legends

### Table 1. Baseline patient and disease characteristics

BRAF<sup>i</sup>, BRAF inhibitor; LDH, lactate dehydrogenase; ULN, upper limit of the normal range; WHO, World Health Organization.

\*Includes 2 patients in the stepwise group; 300 mg (first 15 days of treatment) to 450 mg (from day 15 onward if patients experienced no clinically significant grade 3 or 4 adverse events at the lower dose level).

### Table 2. Adverse events occurring in ≥30% of patients in the dose-expansion phase

PPED, palmar-plantar erythrodysesthesia.

\*Includes 2 patients in the stepwise group; 300 mg (first 15 days of treatment) to 450 mg (from day 15 onward if patients experienced no clinically significant grade 3 or 4 adverse events at the lower dose level).

### Table 3. Summary of best overall response for patients in the dose-escalation and dose-expansion phases

CR, complete response; ORR, overall response rate; PR, partial response.

\*Includes 2 patients in the stepwise group; 300 mg (first 15 days of treatment) to 450 mg (from day 15 onward if patients experienced no clinically significant grade 3 or 4 adverse events at the lower dose level).

<sup>†</sup>Unconfirmed CR due to no confirmatory scan before data cutoff.

<sup>‡</sup>Unconfirmed CR/PR due to PR followed by progressive disease at the confirmatory scan.

<sup>§</sup>Unconfirmed CR/PR due to PR followed by patient discontinuation due to AE.

## Figure Legends

### Figure 1. Pharmacologic characterization of encorafenib

(A) Impact of BRAF inhibitors on phosphorylated MEK (pMEK) in *BRAF* V600E-mutant cells; (B) Tumor pERK inhibition and plasma encorafenib concentration across varying doses (0.6-60 mg/kg) of encorafenib. Female nude mice bearing A375 (*BRAF* V600E) human melanoma tumor xenografts were given a single oral dose of vehicle or encorafenib, and plasma and tumor samples were collected at predetermined time points. Phosphorylated ERK levels were measured in tumor lysates using a Meso Scale Discovery assay (normalized to total ERK) and the percent of inhibition was plotted relative to vehicle controls; (C) Efficacy of encorafenib, dabrafenib, and vemurafenib in the A375 human melanoma tumor xenograft model grown in nude mice (n=5 per group). Encorafenib and vemurafenib were dosed twice daily on an 8/16 hour interval and dabrafenib was dosed QD by oral gavage for 14 days. Tumor volume and body weight were measured twice per week. No signs of toxicity or mortality were observed (see Supplementary Figure S2A); (D) Efficacy of encorafenib in the HMEX1906 primary human melanoma xenograft model grown in nude mice (n=8 per group); (E) Waterfall plot of change in volume for each tumor in each treatment group from the experiment depicted in 1D. Each bar represents the maximum treatment effect observed for each mouse; (F) Mice from the experiment depicted in 1D were dosed continually, and the endpoint of

conditional survival was marked on the day at which each tumor regrew to its predose baseline.

BID, twice daily; QD, once daily; RFU, relative fluorescence units.


**Figure 2. Median semilogarithmic concentration-time profiles for plasma concentration of encorafenib by time and treatment group in the dose-escalation phase**

cps, capsule; m, microemulsion; QD, once daily.

**Figure 3. Efficacy of encorafenib by treatment group**


Best percentage change in sum of longest diameters in target lesion from baseline by treatment group in the dose-escalation (A) and dose-expansion phases (B); Kaplan-Meier plot of progression-free survival (C) and overall survival (D) by patient group in the dose-escalation phase.

**Figure 1**




**Figure 2**

**Cycle 1 Day 1**



**Cycle 1 Day 15**




Downloaded from [clincancerres.aacrjournals.org/](https://clincancerres.aacrjournals.org/) on August 28, 2017. © 2017 American Association for Cancer Research.

Encofenib 50 mg (cvs - QD) Encofenib 50 mg (m - QD) Encofenib 100 mg (cvs - QD)  
Encofenib 100 mg (m - QD) Encofenib 150 mg (cvs - QD) Encofenib 200 mg (cvs - QD)  
Encofenib 700 mg (cvs - QD)


**Figure 3**

**A**

**BRAFi-naïve**




**BRAFi-pretreated**



**Treatment group**

○ 50 mg QD    △ 150 mg QD    ◇ 200 mg QD    □ 300 mg QD    ▽ 450 mg QD    × 700 mg QD  
● 100 mg QD    ▲ 75 mg BID    ◆ 100 mg BID    ■ 150 mg BID    ▴ 550 mg QD

**B**



**Treatment group**


○ BRAFi-naïve 300 mg QD    △ BRAFi-pretreated 300 mg QD    ▲ Step-wise group  
● BRAFi-naïve 450 mg QD    ▲ BRAFi-pretreated 450 mg QD

**C**

**Median progression-free survival, months**

BRAFi-naïve: 12.35 [7.43; NA]

BRAFi-pretreated: 1.91 [0.92; 3.68]




**D**

**Median overall survival, months**

BRAFi-naïve: NA

BRAFi-pretreated: 9.07 [3.68; 10.84]



# Clinical Cancer Research

## Phase I Dose-Escalation and -Expansion Study of the BRAF Inhibitor Encorafenib (LGX818) in Metastatic BRAF-Mutant Melanoma

Jean-Pierre Delord, Caroline Robert, Marta Nyakas, et al.

*Clin Cancer Res* Published OnlineFirst June 13, 2017.

<b>Updated version</b>	Access the most recent version of this article at: doi: <a href="https://doi.org/10.1158/1078-0432.CCR-16-2923">10.1158/1078-0432.CCR-16-2923</a>
<b>Supplementary Material</b>	Access the most recent supplemental material at: <a href="http://clincancerres.aacrjournals.org/content/suppl/2017/06/13/1078-0432.CCR-16-2923.DC1">http://clincancerres.aacrjournals.org/content/suppl/2017/06/13/1078-0432.CCR-16-2923.DC1</a>
<b>Author Manuscript</b>	Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

<b>E-mail alerts</b>	<a href="#">Sign up to receive free email-alerts</a> related to this article or journal.
<b>Reprints and Subscriptions</b>	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at <a href="mailto:pubs@aacr.org">pubs@aacr.org</a> .
<b>Permissions</b>	To request permission to re-use all or part of this article, contact the AACR Publications Department at <a href="mailto:permissions@aacr.org">permissions@aacr.org</a> .